
## Steps to Put On personal protective equipment (PPE)

- Always put on essential required PPE when handling either a suspected, probable or confirmed case of viral haemorragic fever.
- The dressing and undressing of PPE should be supervised by another trained member of the team.
- 3 Gather all the necessary items of PPE beforehand. Put on the scrub suit in the changing room.



4 Put on rubber boots. If not available, make sure you have closed, puncture and fluid resistant shoes and put on overshoes.



OR, IF BOOTS UNAVAILABLE



5 Place the impermeable gown over the scrubs.



- 6 Put on face protection:
- 6a Put on a medical mask.



6b Put on goggles or a face shield.





If available, put a head cover on at this time.



8 Perform hand hygiene.



9 Put on gloves\* (over cuff).



10 If an impermeable gown is not available, place waterproof apron over gown.



## While wearing PPE:

- Avoid touching or adjusting PPE
- · Change gloves between patients
- Remove gloves if they become torn or damaged
- Perform hand hygiene before putting on new gloves
- \* Use *double gloves* if any strenuous activity (e.g. carrying a patient or handling a dead body) or tasks in which contact with blood and body fluids are anticipated. Use *heavy duty/rubber gloves* for environmental cleaning and waste management.



## Steps to remove personal protective equipment (PPE)

1 Remove waterproof apron and dispose of safely. If the apron is to be reused, place it in a container with disinfectant.




2 If wearing overshoes, remove them with your gloves still on (If wearing rubber boots, see step 4).



3 Remove gown and gloves and roll inside-out and dispose of safely.



4 If wearing rubber boots, remove them (ideally using the boot remover) without touching them with your hands. Place them in a container with disinfectant.



5 Perform hand hygiene.



If wearing a head cover, remove it now (from behind the head).



- 7 Remove face protection:
- 7a Remove face shield or goggles (from behind the head). Place eye protection in a separate container for reprocessing.



7b Remove mask from behind the head. When removing mask, untie the bottom string first and the top string next.



8 Perform hand hygiene.



**Source:** Modified from Clinical Management of Patients with Viral Haemorrhagic Fever: A pocket Guide for the Front-line Health Worker. World Health Organization, 2014

